**JBP****Journal of Biochemicals and Phytomedicine**

eISSN: 2958-8561



# Ethnobotanical Study of Effective Medicinal Plants as Laxative (Anti-constipation): A Case Study in Arasbaran Forests, Northwest Iran

Ehsan Bahmani<sup>1</sup> , Fariba Bahmani<sup>2</sup> , Saber Abbaszadeh<sup>3\*</sup> <sup>1</sup>Faculty of Agriculture and Natural Resources, Razi University, Kermanshah, Iran<sup>2</sup>Ilam University of Medical Sciences, Ilam, Iran<sup>3</sup>Department of Biochemistry and Genetics, School of Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran**ARTICLE INFO***Article Type:*

Research

*Article History:*

Received: 19 Nov 2022

Revised: 10 Jun 2023

Accepted: 10 Jun 2023

Available online: 30 Jun 2023

*Keywords:*Ethnobotany,  
Medicinal plants,  
Laxative,  
Anti-constipation agents,  
Constipation*\*Corresponding author:*E-mail: [saberabaszade1370@gmail.com](mailto:saberabaszade1370@gmail.com)**ABSTRACT**

**Introduction:** Constipation is recognized as a common gastrointestinal problem that many sick and healthy individuals suffer from. Laxative therapy is a reliable way to treat constipation. Several medicinal plants exert laxative potential. Hence, we decided to identify medicinal plants with laxative properties in Arasbaran Forests, East Azerbaijan Province, Northwest Iran.

**Methods:** This ethnobotanical study was conducted in Arasbaran Forests, East Azerbaijan Province, and Northwest Iran from October 2022 to December 2022. Questionnaires were distributed among 20 traditional medicine doctors and finally, traditional information on the therapeutic effect of medicinal plants was collected through interviews. Demographic information from indigenous people who have traditional botanical knowledge was also collected. Eventually, the data were analyzed.

**Results:** Our work revealed that thirteen medicinal herbs from 11 herbal families have potent laxative effects. The Berberidaceae, Equisetaceae, Caprifoliaceae, Celastereae, Asteraceae, Cucurbitaceae, Euphorbiaceae, Grossulariaceae, Polygonaceae, Polypodiaceae, and Rosaceae families had laxative effects. Fruit, leaves, aerial parts, bark, root, and rhizome were the most effective parts, respectively. These medicinal plants were administered in decoction and fresh form.

**Conclusion:** The findings show the traditional laxative effects of medicinal plants against constipation in this region, which have traditional uses. Therefore, identifying the phytochemical constituents of these plants and understanding the mechanism of their laxative effect can be helpful.

*Please cite this paper as:*Bahmani E, Bahmani F, Abbaszadeh S. Ethnobotanical Study of Effective Medicinal Plants as Laxative (Anti-constipation): A Case Study in Arasbaran Forests, Northwest Iran. *Journal of Biochemicals and Phytomedicine*. 2023; 2(2): 59–63. doi: 10.34172/jbp.2023.12.

## Intorduction

Constipation is a common complication that a wide range of healthy and sick people suffer from (Daniali et al., 2020). It has been estimated that constipation has a high prevalence equal to 16% all over the world. However, statistics from Western industrial countries show prevalence of constipation near 30% (Salari et al., 2023; Klaschik et al., 2003). Constipation is a phenomenon that is experienced by the majority of people throughout their lives at least one time, regardless of age or health status (Salari et al., 2023). There are many causes, including diseases, lifestyle and the use of drugs such as narcotics, in the occurrence of constipation (Klaschik et al., 2003; Xu et al., 2021). Constipation could be an emergency condition that may lead to serious gastrointestinal complications such as hemorrhoids, anal fissure, rectal bleeding, and rectal prolapse eventually (Faghihi et al. 2022). Constipation is an unpleasant condition that needs appropriate assessment and immediate treatment. Constipation treatment is a multi-method process that requires changes in behavior, lifestyle, nutritional status and drug therapy with laxatives for faster and more stable treatment (Camilleri et al., 2017). Changing the behavior, lifestyle and nutritional status of a person with constipation through increasing physical activity, increasing fluid intake, increasing consumption of vegetables and fiber-rich foods are effective approaches in managing constipation (Klaschik et al., 2003). Laxative therapy is the selective pharmacotherapy of constipation. Bulk-forming laxatives, stimulant laxatives, osmotic laxatives, and lubricating agents are most common used laxatives agents to treat constipation (Bassotti et al., 2023). The numerous side effects of synthetic laxatives have increased the desire of researchers to use natural alternatives (Khan et al., 2020). Previous studies have demonstrated the beneficial effects of medicinal plants on constipation. Medicinal plants are well known for their various therapeutic effects including antioxidant, antimicrobial and anti-inflammatory properties (Sulaiman et al., 2022; Amiri et al., 2023; Zangeneh et al., 2023; Baharvand Ahmadi et al., 2023; khajehpour A, Javadian, 2020). In addition to the use of medicinal plants as remedies, they have been recognized as a reliable food sources. Medicinal herbs can improve health status because of their phytochemical constituents. For instance, it has been reported that medicinal plants relieve constipation because they are rich in fiber (Sharma et al., 2021; Jasim et al. 2023; Eftekhari et al., 2022; Baniesmaeili et al. 2023). Hence, the aim of this study was to identify effective medicinal plants as laxative in Arasbaran Forests, East Azerbaijan Province, and Northwest Iran.

## Materials and Methods

### Area of Study

The present ethnobotanical study was performed in Arasbaran Forests, East Azerbaijan Province, Northwest Iran.


### Data Collection

The current study was a cross-sectional systematic and ethnobotanical research that was conducted in the


period from February 2021 to May 2021. First, standard questionnaires were prepared and these questionnaires were administered to 20 traditional medicine doctors (five women and 15 men). The questionnaire was distributed among traditional medicine doctors. The results obtained from the traditional information of the studied subjects were recorded directly in the questionnaire tables and finally the data were analyzed using Excel software (Baharvand Ahmadi et al., 2015).

## Results

Our findings highlighted that 13 medicinal herbs from eleven families were administered as laxative (Table 1). It has also been shown that the medicinal herbs in Berberidaceae, Equisetaceae, Caprifoliaceae, Celasteraceae, Asteraceae, Cucurbitaceae, Euphorbiaceae, Grossulariaceae, Polygonacea, Polypodiaceae, and Rosaceae families were laxative (Table 2). Moreover, it has been understood that fruit (41%), leaves (23%), aerial parts (6%), bark (6%), root (18%), and rhizome (6%) were the most usable part of medicinal plants (Figure 1). In addition, it has been observed that decoction (77%) and infusion (23%) were the most important types of preparation methods of the medicinal plants, respectively (Figure 2).



**Figure 1.** The percentage of plant organs used traditionally as laxative in Arasbaran forests



**Figure 2.** The percentage of traditional used forms of medicinal plants in Arasbaran forests as laxative

**Table 1.** Medicinal plants used as laxatives in Arasbaran, Iran

Scientific Name	Family	Persian Name	Usable Part of Plant	Preparation Methods	N	%
<i>Berberis vulgaris</i> L.	Berberidaceae	Zereshk	Fruit & leave	Decoction	8	53.33
<i>Equisetum arvense</i> L.	Equisetaceae	Dome asb	Aerial parts	Decoction	7	46.66
<i>Sambucus nigra</i> L.	Caprifoliaceae	Angoore koli	Fruit	Fresh	9	60
<i>Sambucus ebulus</i>	Caprifoliaceae	Aghti	Bark	-	4	26.66
<i>Euonymus latifolius</i> (L.) Mill.	Celasteraceae	Gooshvarak	Fruit & leave	Decoction	3	20
<i>Cichorium intybus</i> L.	Asteraceae	Kasni	Root & leave	Decoction	4	26.66
<i>Ecballium elaterium</i> A. Rich	Cucurbitaceae	Khiareh vahshi	Root & fruit	Decoction	6	40
<i>Euphorbia helioscopia</i> L.	Euphorbiaceae	Ferfion	Root	Decoction	5	33.33
<i>Ribes orientale</i> Desf	Grossulariaceae	Angoore sharghi	Fruit	Decoction	8	53.33
<i>Rumex acetosa</i>	Polygonacea	Torshak	Leave	Fresh	2	13.33
<i>Polypodium vulgare</i>	Polypodiaceae	Befaiej	Rhizome	Decoction	7	46.66
<i>Prunus spinosa</i> L.	Rosaceae	Gojeh vahshi	Fruit	Fresh & Decoction	4	26.66
<i>Sorbus boissieri</i> C. K. syn. <i>S. aucuparia</i> L.	Rosaceae	Tis	Fruit	Decoction	1	6.66

N: The number of people who stated in this study that this plant has laxative effects; %: The percentage of people who stated in this study that this plant has laxative effects; -: Not announced.

**Table 2.** Distribution of medicinal plants as laxative in Arasbaran forests on this study

Row	Herbal family	Number
1	Berberidaceae	1
2	Equisetaceae	1
3	Caprifoliaceae	2
4	Celasteraceae	1
5	Asteraceae	1
6	Cucurbitaceae	1
7	Euphorbiaceae	1
8	Grossulariaceae	1
9	Polygonacea	1
10	Polypodiaceae	1
11	Rosaceae	2

## Discussion

Numerous studies have documented the role of medicinal herbs as laxative to treat constipation (Akram et al., 2022). In Iranian traditional medicine, it is recommended to use herbs as laxatives to solve the problem of constipation (Karami et al., 2020). Therefore, the present ethnobotanical study that we know was the first to investigate the effect of medicinal plants of Arasbaran forests as a laxative on constipation. The results of the present ethnobotanical study revealed that thirteen medicinal herbs from eleven families including *Berberis vulgaris* L. known as barberry, *Equisetum arvense* L. (field horsetail), *Sambucus nigra* L. known as elder, *Sambucus ebulus*, *Euonymus latifolius* (L.) Mill., *Cichorium intybus* L. known as chicory, *Ecballium elaterium* A. Rich (squirting cucumber), *Euphorbia helioscopia* L., *Ribes orientale* Desf, *Rumex acetosa* known as Sorrel, *Polypodium vulgare* (common polypody), *Prunus spinosa* L. known as blackthorn and *Sorbus boissieri* C. K. syn. *S. aucuparia* L. were prescribed as laxative. In addition, the findings of the current study highlighted that medicinal plants in Berberidaceae, Equisetaceae, Caprifoliaceae, Celasteraceae, Asteraceae, Cucurbitaceae, Euphorbiaceae, Grossulariaceae, Polygonacea, Polypodiaceae, and Rosaceae families were recognized as potent laxative, respectively. Our work also demonstrated that fruit, leaves, aerial parts,

bark, root, and rhizome were the most usable part of the mentioned plants. Furthermore, we found that these medicinal plants were administrated in the form of decoction and infusion. In recent decades, researchers have focused on the use of natural agents to treat various diseases. Herbal plants have a long history in the treatment of gastrointestinal diseases. In fact, medicinal plants with various phytochemical compounds, including flavonoids, phenolic compounds, terpenoids (or isoprenoids), saponins, and vitamins cause multiple therapeutic properties such as antioxidant, antimicrobial, anticancer and anti-inflammatory properties. Medicinal plants, with their anti-inflammatory and antimicrobial properties, deal with inflammation and pathogenic microbes in the digestive system, which are the cause of many diseases in these organs. On the other hand, the presence of high amounts of plant fibers in them increases the volume of feces, increases intestinal movements, and thus prevents constipation (Karami et al., 2020; Hashem et al., 2015). In a similar ethnobotanical study conducted by Karami N et al., the effectiveness of medicinal plants for treatment of constipation was evaluated in Shahrekord city, Chaharmahal & Bakhtiari province, Iran. The results of their study indicated that sixteen medicinal plants including flaxseed (*Descurainia sophia* (L.) Prantle.),

fiddle dock (*Rumex pulcher* L.), *Alyssum* spp. Stead. Ex Boiss., almond (*Prunus amygdalus* L.), Persian manna (*Astragalus adscondens* Bioss. & Hauskn), *Echinops persicus* Stev. & Fisch., broadleaf plantain (*Plantago major* L.), *Rheum ribes* L., pink cotton lamb's ear (*Stachys lavandulifolia*), *Plantago major*, *Alcea* spp., licorice (*Glycyrrhiza glabra* L.), wild rue (*Peganum harmala* L.), *Pistacia atlantica* Desf. and common basilisk (*Prangos ferulacea* L.) showed a potent anti-constipation effect (Karami et al., 2020). The results of this study in line with the results of our study documented the therapeutic effects of medicinal plants on constipation as laxative. In another similar study, Hashem Dabaghian F et al. reported that Iranian traditional medicinal plants such as golden shower tree (*Cassia fistula*), *Cotoneaster nummularia* Fisch., tamarind (*Tamarindus indica*), camelthorn (*Alhagi camelorum*), and Damask rose (*Rosa damascene*) and almond oil were used to treat constipation especially during pregnancy (Hashem et al., 2015).

## Conclusions

Our findings revealed that medicinal plants used in Arasbaran have potent therapeutic effect on constipation as laxative. The chemical compounds found in medicinal plants are the main factor in helping to treat constipation by increasing bowel movements and volume. Therefore, studying more about the role of herbal compounds in improving digestive problems such as constipation can be a factor in reducing the side effects of chemical drugs.

## Declarations

### Conflict of interest

There is no conflict of interest.

## Acknowledgement

None.

## Consent for publications

The author approved the manuscript for publication.

## Funding/support

None.

## Authors' contributions

SA and EB conceived the research idea. FB and SA designed the work. SA carried out the experiment. FB and EB wrote the first draft of the manuscript. SA carried out the literature search. EB carried out the statistical analysis, and SA supervised the study. All authors read and approved the final manuscript for publication.

## Ethical considerations

Ethical issues (including plagiarism, misconduct, data fabrication, falsification, double publication or submission, redundancy) have been completely observed by the author.

## References

- Akram M, Thiruvengadam M, Zainab R, Daniyal M, Bankole MM, Rebezov M, et al. Herbal medicine for the management of laxative activity. *Current Pharmaceutical Biotechnology*. 2022;23(10):1269-83. <https://doi.org/10.2174/1389201022666210812121328>.
- Amiri MM, Garnida Y, Almulla AF, Abduljabbar AS, Jalil AT, Mazaheri Y, Ebrahimi Y, Shariatifar N. Herbal therapy for hemorrhoids: An overview of medicinal plants affecting hemorrhoids. *Advancements in Life Sciences*. 2023; 10(1):22-8.
- Baharvand Ahmadi B, Khajoei Nejad F, Papi S, Eftekhari Z. Phytotherapy for heart tonic: An ethnobotanical study in Dehloran City, Ilam Province, Western Iran. *Caspian Journal of Environmental Sciences*. 2023; 1-5. doi: 10.22124/cjes.2023.6192
- Baharvand-Ahmadi B, Bahmani M, Tajeddini P, Rafieian-Kopaei M, Naghdi N. An ethnobotanical study of medicinal plants administered for the treatment of hypertension. *Journal of Renal Injury Prevention*. 2016;5(3):123. <https://doi.org/10.15171/jrip.2016.26>.
- Baniesmaeili SY, Akbarzadeh A, Riazi G, Abdollahi F, Niromand M. Effects of dietary Moringa oleifera leaf powder and ethanolic extract on expression function of immune genes of *Litopenaeus vannamei*. *Aquatic Animals Nutrition*. 2023; 9(1): 27-41. doi: 10.22124/janb.2023.24268.1201
- Bassotti G, Battaglia E. Considerations for laxatives in terms of their interactions with other drugs. *Expert Opinion on Drug Metabolism & Toxicology*. 2023; 19(3): 121-3. <https://doi.org/10.1080/17425255.2023.2193326>.
- Camilleri M, Ford AC, Mawe GM, Dinning PG, Rao SS, Chey WD, et al. Chronic constipation. *Nature Reviews Disease Primers*. 2017;3(1):1-19. <https://doi.org/10.1038/nrdp.2017.95>.
- Daniali M, Nikfar S, Abdollahi M. An overview of interventions for constipation in adults. *Expert Review of Gastroenterology & Hepatology*. 2020;14(8):721-32. <https://doi.org/10.1080/17474124.2020.1781617>.
- Eftekhari Z, Patra I, Hamza TA, Adhab AH, Hachim SK, Abulkassim R, Hameed NM, Naghdi N. Evaluation of the total antioxidant capacity of bitter and sweet varieties of *Ferula assa-foetida* and *Bunium persicum*. *Advancements in Life Sciences*. 2022;9(3):363-7.
- Faghihi A, Zohalinezhad ME, Najafi Kalyani M. Comparison of the effects of abdominal massage and oral administration of sweet almond oil on constipation and quality of life among elderly individuals: A single-blind clinical trial. *BioMed Research International*. 2022;2022. <https://doi.org/10.1155/2022/9661939>.
- Hashem Dabaghian F, Taghavi Shirazi M, Amini Behbahani F, Shojaee A. Interventions of Iranian traditional medicine for constipation during pregnancy. *Journal of Medicinal Plants*. 2015;14(53):58-68. <http://dorl.net/dor/20.1001.1.2717204.2015.14.53.7.8>.

Jasim SA, Abdelbasset WK, Jawad MA, Bokov, DO, Thangavelu L, Manouchehri A. Tramadol toxicity phytotherapy: The protective role of medicinal plants against tramadol toxicity. *Caspian Journal of Environmental Sciences*. 2023; 21(1): 227-243. doi: 10.22124/cjes.2023.6234

Karami N, Karimi M, Bahmani M. Constipation: An ethno-botanical study of medicinal plants used for constipation in Shahrekord city, Chaharmahal & Bakhtiari province, Iran. *Plant Biotechnol Persa*. 2020;2(1):1-7. <http://dx.doi.org/10.29252/pbp.2.1.1>.

Khajehpour A, Javadian R. Effects of orally-administered *Echinacea purpurea* on growth indices, hematology and innate immunity in Siberian sturgeon (*Acipenser baerii*). *Aquatic Animals Nutrition*. 2020; 6(1): 13-25. doi: 10.22124/janb.2020.17007.1094

Khan S, Khan SU. Adverse drug event of hypokalaemia-induced cardiotoxicity secondary to the use of laxatives: A systematic review of case reports. *Journal of Clinical Pharmacy and Therapeutics*. 2020;45(5):927-36. <https://doi.org/10.1111/jcpt.13204>.

Klaschik E, Nauck F, Ostgathe C. Constipation—modern laxative therapy. *Supportive Care in Cancer*. 2003;11:679-85. <https://doi.org/10.1007/s00520-003-0525-x>.

Salari N, Ghasemianrad M, Ammari-Allahyari M, Rasoulpoor S, Shohaimi S, Mohammadi M. Global prevalence of constipation in older adults: a systematic review and meta-analysis. *Wiener Klinische Wochenschrift*. 2023; 135:389-98. <https://doi.org/10.1007/s00508-023-02156-w>.

Sharma A, Khanna S, Kaur G, Singh I. Medicinal plants and their components for wound healing applications. *Future Journal of Pharmaceutical Sciences*. 2021;7(1):1-13. <https://doi.org/10.1186/s43094-021-00202-w>.

Sulaiman AN, Arzai AH, Taura DW. Ethnobotanical survey: A comprehensive review of medicinal plants used in treatment of gastro intestinal diseases in Kano state, Nigeria. *Phytomedicine Plus*. 2022;2(1):100180. <https://doi.org/10.1016/j.phyplu.2021.100180>.

Xu Y, Amdanee N, Zhang X. Antipsychotic-induced constipation: a review of the pathogenesis, clinical diagnosis, and treatment. *CNS drugs*. 2021;35(12):1265-74. <https://doi.org/10.1007/s40263-021-00859-0>.

Zangeneh MM, Zangeneh A, Etemadi N, Tabatabaei Aghda SJ, Pourkamalzadeh M, Zand S, et al. The effect of nano silver particles of the aqueous extract of *Artemisia aucheri* on the healing process of skin wounds in rats. *Plant Biotechnology Persa*. 2023; 5(1):1-7.