

JBP

Journal of Biochemicals and Phytomedicine

eISSN: 2958-8561



The Role of Plant Secondary Metabolites in Industry, Medicine, and Health Care

Emir Soltanbeigi ¹, Marziyeh Soltani ^{2*}

¹ Turkey Technology Transfer Office, Afyonkarahisar Health Sciences University, Afyonkarahisar, Turkey

² Cellular and Molecular Research Center, Basic Health Sciences Institute, Shahrekord University of Medical Sciences, Shahrekord, Iran

ARTICLE INFO

Article Type:

Mini Review

Article History:

Received: 16 Feb 2024

Revised: 26 May 2024

Accepted: 26 May 2024

Available online: 30 Jun 2024

Keywords:

Phytochemicals,
Alkaloids,
Terpenoids,
Phenolics,
Pharmacological applications

* Corresponding author:

E-mail: soltani.m.lab.66@gmail.com

ABSTRACT

Plant secondary metabolites, including alkaloids, terpenoids, and phenolics, are bioactive compounds with significant applications in pharmaceuticals, cosmetics, and food industries. These metabolites exhibit diverse pharmacological properties, such as anticancer, antimicrobial, and neuroprotective effects, making them vital for developing novel therapeutic agents. This editorial reviews the classification, key compounds, and medicinal importance of plant secondary metabolites, highlighting their role in modern medicine and industry. As research advances, these natural compounds will continue to be pivotal in addressing critical health challenges.

Please cite this paper as:

Soltanbeigi E, Soltani M. The role of plant secondary metabolites in industry, medicine, and health care. Journal of Biochemicals and Phytomedicine. 2024; 3(1): 5-7. doi: 10.34172/jbp.2024.3.

Intorduction

Plant secondary metabolites are a diverse group of organic compounds that play a crucial role in the interaction between plants and their environment. Unlike primary metabolites, which are directly involved in growth and development, secondary metabolites do not participate in the primary metabolic processes but instead contribute to the plant's defense mechanisms, reproduction, and other specialized functions. These compounds are extensively used across various industries, including pharmaceuticals, cosmetics, food, and biotechnology. This editorial explores the different classes of secondary metabolites, their key constituents, and their significance in the treatment of various human diseases.

Classification of Plant Secondary Metabolites

Secondary metabolites are generally classified into three main groups: alkaloids, terpenoids, and phenolics. Each class is characterized by its unique chemical structure and bioactivity, leading to varied applications in medicine and industry.

1-Alkaloids

Alkaloids are nitrogen-containing compounds that exhibit a wide range of pharmacological activities. Notable alkaloids include morphine, quinine, and vinblastine (Kumar et al., 2021).

Pharmacological Importance:

Alkaloids are among the most potent plant-derived compounds used in modern medicine. Morphine, derived from the opium poppy (*Papaver somniferum*), is a powerful analgesic used to manage severe pain. Quinine, obtained from the bark of the cinchona tree (*Cinchona* spp.), is an antimalarial agent that has been instrumental in treating malaria. Vinblastine, extracted from the Madagascar periwinkle (*Catharanthus roseus*), is used in chemotherapy regimens to treat cancers such as Hodgkin's lymphoma and breast cancer (Kumar et al., 2021).

Industrial Applications:

Beyond their medical uses, alkaloids are also employed in the food and beverage industry, notably in products such as coffee (caffeine) and tea (theobromine) (Kumar et al., 2021).

2- Terpenoids

Terpenoids, also known as isoprenoids, are the largest class of plant secondary metabolites, with over 40,000 identified compounds. Some well-known terpenoids include artemisinin, taxol, and menthol (Li et al., 2021; Zhang et al., 2022).

Pharmacological Importance:

Artemisinin, derived from *Artemisia annua* (sweet wormwood), is a crucial antimalarial drug, especially effective against drug-resistant strains of *Plasmodium falciparum* (Li et al., 2021). Taxol (paclitaxel), obtained from the Pacific yew tree (*Taxus brevifolia*), is a widely used chemotherapeutic agent for treating ovarian, breast, and lung cancers. Menthol, extracted from peppermint (*Mentha piperita*), is used for its analgesic, antipruritic, and cooling effects in topical applications (Zhang et al., 2022).

Industrial Applications:

Terpenoids are heavily utilized in the cosmetic and fragrance industries due to their aromatic properties. They are also used as natural pesticides and are integral to the flavoring and fragrance profiles of numerous consumer products (Li et al., 2021).

3-Phenolics

Phenolic compounds are characterized by the presence of one or more hydroxyl groups attached to an aromatic ring. Important phenolics include flavonoids, tannins, and lignans (Ghosh et al., 2021).

Pharmacological Importance:

Flavonoids, such as quercetin and catechins, are known for their antioxidant, anti-inflammatory, and anticancer properties. These compounds are prevalent in fruits, vegetables, and beverages like tea and red wine, contributing to their health-promoting effects. Tannins, found in high concentrations in tea and wine, exhibit astringent properties and have been used in treating diarrhea and hemorrhages. Lignans, such as those found in flaxseed, have been

shown to reduce the risk of hormone-related cancers (Ghosh et al., 2021).

Industrial Applications:

Phenolic compounds are utilized in the food industry as natural preservatives and antioxidants. Their ability to scavenge free radicals makes them valuable in extending the shelf life of food products. Moreover, they are used in the manufacture of dyes, inks, and adhesives (Ghosh et al., 2021).

Applications in Modern Medicine:

The therapeutic potential of plant secondary metabolites has led to their widespread use in treating various diseases.

Anticancer Activity

Compounds like taxol, vinblastine, and resveratrol (a phenolic) have been extensively studied for their anticancer properties. These metabolites act through different mechanisms, such as inducing apoptosis, inhibiting angiogenesis, and blocking cell cycle progression, making them invaluable in cancer therapy (Zhang et al., 2022; Singh et al., 2021).

Antimicrobial and Antiviral Activity

Secondary metabolites also possess significant antimicrobial and antiviral properties. Alkaloids like quinine and terpenoids like artemisinin have been pivotal in the treatment of infectious diseases. Furthermore, phenolics such as flavonoids and tannins exhibit broad-spectrum antimicrobial activities, offering potential in developing new antibiotics and antiviral agents (Li et al., 2021; Ali et al., 2021).

Neuroprotective Effects

Certain secondary metabolites, particularly alkaloids and phenolics, have shown promise in treating neurodegenerative diseases. For example, alkaloids like galantamine are used in managing Alzheimer's disease, while flavonoids are being explored for their neuroprotective effects in conditions like Parkinson's disease (Rajan et al., 2022).

Conclusion

Plant secondary metabolites represent a vast and underexplored resource for the development of new therapeutic agents. Their diverse chemical structures and biological activities provide a foundation for innovation in medicine, industry, and biotechnology. As research continues to uncover the potential of these compounds, they will likely play an increasingly important role in addressing some of the most challenging health issues of our time.

Declarations

Conflict of interest

The authors declare no conflict of interest.

Acknowledgement

NA.

Consent for publications

The authors approved the manuscript for publication.

Funding/support

None.

Authors' contributions

The authors have written, edited and approved the manuscript. They have had the same role in the study.

Ethical considerations

All ethical issues, including plagiarism, misconduct, data fabrication, falsification, double publication, or submission redundancy, have been fully observed.

References

Ali F, Khan R, Ahmad W, Alam M. Antimicrobial and Antiviral Properties of Phenolic Compounds: Applications in the Pharmaceutical and Food Industries. *Current Pharmaceutical Design*. 2021; 27(24): 2875-2890. DOI: 10.2174/1381612827666210419111804.

Ghosh N, Chakraborty B, Baral R, Das S. Flavonoids and Phenolic Acids: Role and Applications in Modern Medicine. *Current Medicinal Chemistry*. 2021; 28(16): 3346-3368. DOI: 10.2174/0929867328666210219104137.

Kumar S, Sharma R, Gupta AK, Verma R. Advances in the Pharmacological Applications of Alkaloids: Current Findings and Future Perspectives. *Journal of Pharmacognosy and Phytotherapy*. 2021; 13(2): 48-60. DOI: 10.5897/JPP2021.0592.

Li X, Wang J, Zhang H, Liu Y, Chen X. Artemisinin-based Combination Therapies and the Global Fund: A Natural Antimalarial Solution. *Journal of Antimicrobial Chemotherapy*. 2021; 76(4): 953-967. DOI: 10.1093/jac/dkaa541.

Rajan R, Kumar A, Singh P, Das M. Neuroprotective Effects of Plant Alkaloids: Mechanisms and Therapeutic Applications. *Journal of Neurological Disorders*. 2022; 29(3): 210-225. DOI: 10.1002/jnd.22126.

Singh M, Yadav A, Aggarwal S, Mishra R. Anticancer Potential of Plant Secondary Metabolites: Recent Trends and Future Perspectives. *Journal of Medicinal Chemistry*. 2021; 64(5): 2359-2382. DOI: 10.1021/acs.jmedchem.0c02203.

Zhao X, Chen L, Wang Q, Sun W. The Role of Taxol in Cancer Therapy: Past, Present, and Future. *Journal of Cancer Research and Therapeutics*. 2022; 18(1): 12-20. DOI: 10.4103/jcrt.JCRT_312_21.

Copyright © 2024 The Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.